

A New Class of Generalized U-Birecurrent Finsler Spaces and Their Curvature Properties

Kamal Salem Naji Nasr¹

Adel Mohammed Ali Al-Qashbari²

Dept. of Math's., Faculty of Educ. Aden, Univ. of Aden, Aden, Yemen

Email: www.kamalsalem1983@gmail.com

²Department of Biomedical Engineering, Faculty of Engineering and Computers, University of Science and Technology, Aden, Yemen, Email: a.alqashbari@ust.edu.

DOI: [https://doi.org/10.47372/jef.\(2026\)20.1.193](https://doi.org/10.47372/jef.(2026)20.1.193)

Abstract: In this paper, we introduce a new class of generalized U-birecurrent Finsler spaces defined in terms of second-order Berwald covariant derivatives of the $h(v)$ -curvature tensor. This new structure extends existing concepts of recurrent and birecurrent Finsler spaces by incorporating higher-order differential conditions. Several fundamental properties of the proposed spaces are established, including relationships between the $h(v)$ -curvature tensor and other important curvature tensors such as Cartan's curvature tensors, Berwald curvature tensor, and the normal projective curvature tensor. Furthermore, a series of theorems and corollaries are derived to characterize the geometric behavior of these spaces under generalized recurrence conditions. The obtained results provide a broader framework for the study of curvature structures in Finsler geometry and contribute to the ongoing development of higher-order geometric analysis.

Keywords: Finsler geometry; generalized birecurrent space; U-birecurrence; Berwald covariant derivative; curvature tensor; $h(v)$ -curvature tensor; Cartan curvature tensor; projective curvature tensor.

1. Introduction

Finsler geometry is regarded as a natural and far-reaching generalization of Riemannian geometry, in which the metric depends on both position and direction. This additional degree of freedom allows Finsler spaces to model a wide range of anisotropic phenomena arising in mathematics and physics. In recent years, significant progress has been made in the study of curvature structures and tensorial invariants in Finsler manifolds, particularly in relation to geometric analysis and modern physical applications [32–36].

Curvature tensors play a central role in understanding the intrinsic geometric properties of Finsler spaces. Various types of curvature tensors, including Cartan, Berwald, and projective curvature tensors, have been extensively studied to describe geometric behavior under different connections and transformations. Recent works have emphasized the importance of higher-order curvature tensors and their associated invariants in developing a deeper understanding of Finsler structures [33–36]. Moreover, extensions of curvature tensors from general relativity and relativistic spacetime geometry have also been investigated, highlighting connections between Finsler geometry and theoretical physics [1–3]. A major direction of research in Finsler geometry is the study of recurrent and generalized recurrent structures. These structures describe spaces in which curvature tensors satisfy specific differential conditions, leading to important geometric and symmetry properties. Several types of recurrence, including recurrent, birecurrent, and trirecurrent Finsler spaces, have been introduced and analyzed using different covariant derivatives such as those defined by Cartan and Berwald connections [4-6, 15]. In particular, generalized recurrence decompositions and tensor identities have been studied to understand the behavior of curvature tensors under various geometric constraints [4-6].

More recently, attention has shifted toward higher-order generalizations of recurrence conditions and their applications. A number of studies have introduced new classes of generalized recurrent Finsler spaces using higher-order covariant derivatives, Lie derivatives, and mixed derivative structures [7-12, 18]. These approaches have led to significant developments in understanding the interaction between different curvature tensors, including Weyl, projective, and conharmonic curvature tensors, within generalized geometric

frameworks [10-12, 16]. Furthermore, decomposition techniques and tensorial analysis have been applied to investigate structural properties of Finsler spaces and their geometric invariants [11, 16].

In addition, recent contributions have explored advanced topics such as generalized trirecurrence, higher-order curvature decompositions, and applications of Finsler geometry in areas such as geometric modeling, mechanics, and computational methods [13, 14, 17, 19, 20]. These developments indicate a growing interest in extending classical recurrence concepts to more general and flexible geometric settings.

Despite these advances, most existing studies are primarily limited to first-order recurrence conditions or specific curvature tensors. The role of higher-order Berwald covariant derivatives in defining generalized birecurrent structures, particularly in relation to the $h(v)$ -curvature tensor, remains insufficiently explored. This gap motivates the need for developing a more comprehensive framework that incorporates higher-order differential structures into generalized recurrence theory.

In this paper, we introduce a new class of generalized U-birecurrent Finsler spaces defined via second-order Berwald covariant derivatives of the $h(v)$ -curvature tensor. This formulation provides a natural extension of existing recurrence structures and allows for a deeper analysis of higher-order curvature behavior. We establish several fundamental properties of these spaces and derive new relationships between the $h(v)$ -curvature tensor and other important curvature tensors, including Cartan, Berwald, and normal projective curvature tensors.

The results obtained in this work generalize several known concepts in Finsler geometry and contribute to the ongoing development of higher-order geometric analysis. Moreover, they provide a unified framework for studying curvature interactions in generalized Finsler spaces and open new directions for future research in both theoretical and applied contexts.

Let F_n be an n -dimensional Finsler space equipped with the metric function $F(x, y)$ satisfying the request conditions

Let us consider a set of quantities $g_{ij}(x, y)$ defined by

$$(1.1) \quad g_{ij}(x, y) = \frac{1}{2} \partial_i \partial_j F^2(x, y).$$

Where g_{ij} is positively homogeneous of degree zero in directional arguments.

$$(1.2) \quad y_i = g_{ij} y^j.$$

The quantities g_{ij} and g^{jk} are related by

$$(1.3) \quad g_{ij} g^{jk} = \delta_i^k = \begin{cases} 1 & \text{if } i = k, \\ 0 & \text{if } i \neq k. \end{cases}$$

The vectors y_i and y_k , related with g_{ik} and g^{jk} by

$$(1.4) \quad \begin{aligned} \text{a) } \delta_k^i y_i &= y_k, \quad \text{b) } \delta_k^i y^k = y^i, \quad \text{c) } \delta_j^i g_{ik} = g_{jk}, \quad \text{d) } \partial_j y^i = \delta_j^i, \\ \text{e) } \delta_j^i g^{jk} &= g^{ik}, \quad \text{f) } \delta_j^i \delta_k^j = \delta_k^i, \quad \text{and} \quad \text{g) } \delta_i^i = n. \end{aligned}$$

$$(1.5) \quad C_{ijk} = \frac{1}{2} \partial_i g_{jk} = \frac{1}{4} \partial_i \partial_j \partial_k F^2,$$

where C_{ijk} is known as (h)hv-torsion tensor, it is positively homogeneous of degree -1 in y^i and symmetric in all its indices.

By using Euler's theorem on homogeneous properties, this tensor satisfies the following identities

$$(1.6) \quad \begin{aligned} \text{a) } C_{ijk} y^i &= C_{kij} y^i = C_{jki} y^i = 0, \quad \text{b) } C_{jk}^i y^j = C_{kj}^i y^j = 0, \\ \text{c) } C_{jk}^i y_i &= 0, \quad \text{and} \quad \text{d) } \delta_j^r C_{rkh} = C_{jkh}. \end{aligned}$$

L. Berwald defined covariant derivative for his connection parameter G_{jk}^i . Thus, Berwald's covariant derivative $\mathcal{B}_k X^i$ of an arbitrary vector field X^i with respect to x^k is given by

$$(1.7) \quad \mathcal{B}_k X^i = \partial_h X^i - (\partial_r X^i) G_k^i + X^r G_{rk}^i.$$

Similar to Cartan's covariant derivative, Berwald's covariant derivative of the metric function F , the unit l^i and the vector y^i vanish identical, i.e.

$$(1.8) \quad a) \mathcal{B}_k F = 0, \quad b) \mathcal{B}_k l^i = 0, \quad \text{and} \quad c) \mathcal{B}_k y^i = 0.$$

The covariant derivative of the metric tensor g_{ij} does not vanish in sense of Berwald, i.e. $\mathcal{B}_k g_{ij} \neq 0$.

The covariant derivative of the metric tensor is given by

$$(1.9) \quad \mathcal{B}_k g_{ij} = -2C_{ijkh} y^h = -2y^h \mathcal{B}_h C_{ijk}.$$

The processes of Berwald's covariant differentiation and the partial differentiation commute according to

$$(1.10) \quad (\partial_k \mathcal{B}_h - \mathcal{B}_h \partial_k) T_j^i = T_j^r G_{khr}^i - T_r^i G_{khj}^r, \quad \text{for an arbitrary tensor field } T_j^i.$$

The tensor K_{rkh}^i called Cartan's fourth curvature tensor which is skew-symmetric in its last two lower indices k and h , i.e.

$$(1.11) \quad K_{jkh}^i = -K_{jhk}^i$$

The associate curvature tensor K_{ijkh} of the curvature tensor K_{jkh}^i is given by

$$(1.12) \quad K_{ijkh} = g_{rj} K_{lkh}^r.$$

The associate curvature tensor K_{ijkh} of the curvature tensor K_{jkh}^i is skew-symmetric in its first two lower indices k and h , i.e.

$$(1.13) \quad K_{ijhk} = -K_{jihk}.$$

The K-Ricci tensor K_{jk} , the curvature scalar K and the deviation tensor K_j^i of the curvature tensor K_{jkh}^i are satisfying the following

$$(1.14) \quad a) K_{jki}^i = K_{jk}, \quad b) g^{jk} K_{jk} = K, \quad c) g^{ik} K_{jk} = K_j^i, \quad \text{and} \quad d) K_k^i g_{jh} = K_{jkh}^i.$$

The curvature tensor K_{jkh}^i satisfies the following relations too

$$(1.15) \quad K_{jkh}^i y^j = H_{kh}^i,$$

$$(1.16) \quad H_{jkh}^i = K_{jkh}^i + y^s (\partial_j K_{skh}^i)$$

The tensor R_{jkh}^i called h-curvature tensor (Cartan's third curvature tensor), this tensor is positively homogeneous of degree -1 in y^i and skew-symmetric in its last two lower indices k and h , i.e.

$$(1.17) \quad R_{jkh}^i = -R_{jhk}^i, \quad \text{and satisfies the relations}$$

$$(1.18) \quad R_{jkh}^i = K_{jkh}^i + C_{jm}^i H_{kh}^m, \quad \text{and}$$

$$(1.19) \quad R_{jkh}^i y^j = H_{kh}^i.$$

The R-Ricci tensor R_{jk} , the deviation tensor R_h^r and curvature scalar R of the curvature tensor R_{jkh}^i are given by

$$(1.20) \quad a) R_{jki}^i = R_{jk}, \quad b) R_{ikh}^r g^{ik} = R_h^r, \quad \text{and} \quad c) g^{jk} R_{jk} = R.$$

In contracting the indices i and h in the equation (1.18), we get

$$(1.21) \quad R_{jk} = K_{jk} + C_{jm}^r H_{kr}^m.$$

Berwald's tensors H_{jkh}^i is skew-symmetric in the last two lower indices, i.e. k and h and positively homogeneous of degree zero. i.e

$$(1.22) \quad H_{jkh}^i = -H_{jhk}^i.$$

Torsion tensor H_{kh}^i of Berwald curvature tensor H_{jkh}^i is given by

$$(1.23) \quad a) H_{jkh}^i y^j = H_{kh}^i \quad \text{and} \quad b) H_{jkh}^i = H_{jk}^i.$$

The tensor U_{jkh}^i denoted the tensor Π_{jkh}^i by U_{jkh}^i and defined by

$$(1.24) \quad U_{jkh}^i = G_{jkh}^i + \frac{1}{(n+1)} (\delta_j^i G_{khr}^r + y^i G_{jkh}^r).$$

Also, the tensor U_{jkh}^i is called h(v)-curvature tensor and it is homogeneous of degree -1 in y^i and symmetric in its last two indices and satisfies the following

$$(1.25) \quad a) U_{jkh}^i = U_{jhk}^i, \quad b) U_{jkh}^h = U_{jhk}^h, \quad c) U_{jkh}^i = U_{jhk}^i, \\ d) U_{jkh}^i y^h = U_{jhk}^i y^h = U_{jk}^i, \quad \text{and} \quad e) U_{jkh}^i y^j = 0.$$

The h(v)-Ricci tensor U_{jk} of the projective connection coefficients satisfies the following:

$$(1.26) \quad a) U_{ikh}^i = U_{kh} \quad , \quad b) U_{jkh}^h = U_{jk} \quad , \quad \text{where} \quad c) U_{kh} = \frac{2}{(n+1)} G_{kh} \quad ,$$

where the tensor G_{jk} is components of the call h(v)-Ricci tensor.

The h(v)-torsion tensor U_{jk}^i satisfies

$$(1.27) \quad a) U_{jk}^i = U_{kj}^i \quad , \quad b) U_{jk}^i = U_{jkh}^i y^h \quad , \quad \text{and} \quad c) U_{jr}^r = C_{jr}^r \quad \text{and} \quad d) U_{jkr}^r = C_{jkr}^r \quad .$$

P.N. Pandey obtained a relation between the normal projective curvature tensor N_{jkh}^i and Berwald curvature tensor H_{jkh}^i as follows:

$$(1.28) \quad N_{jkh}^i = H_{jkh}^i - \frac{1}{(n+1)} y^i \partial_j H_{rkh}^r \quad .$$

The normal projective curvature tensor N_{jkh}^i is skew-symmetric in its last two lower indices, i.e.

$$(1.29) \quad N_{jkh}^i = -N_{jhk}^i \quad , \quad \text{and}$$

$$(1.30) \quad N_{jk} = N_{jkr}^r \quad .$$

U-recurrent Finsler space, U^h -birecurrent Finsler space and U-trirecurrent Finsler space are characterized by

$$(1.30) \quad a) \mathcal{B}_l U_{jkh}^i = \lambda_l U_{ikh}^i \quad , \quad \text{where} \quad U_{ikh}^i \neq 0 \quad , \quad \text{and}$$

$$b) \mathcal{B}_m \mathcal{B}_l U_{jkh}^i = a_{lm} U_{ikh}^i \quad .$$

A Finsler space which the h(v)-curvature tensor U_{ikh}^i satisfies the general U_l -recurrent Finsler space and general U_l -birecurrent Finsler space characterized by the following conditions, respectively [10]

$$(1.31) \quad a) \mathcal{B}_l U_{jkh}^i = \lambda_l U_{jkh}^i + b \mu_l (\delta_k^i g_{jh} - \delta_h^i g_{jk}) \quad , \quad U_{jkh}^i \neq 0 \quad ,$$

$$(1.32) \quad b) \mathcal{B}_m \mathcal{B}_l U_{jkh}^i = a_{lm} U_{jkh}^i + b_{lm} (\delta_k^i g_{jh} - \delta_h^i g_{jk}) - 2 \mu_l y^r \mathcal{B}_r (\delta_k^i C_{jhm} - \delta_h^i C_{jkm}) \quad , \quad \text{where}$$

$w_{lm} = \lambda_{lm} + \lambda_l \lambda_m$ and $v_{lm} = \lambda_l \mu_m + \mu_{lm}$ are non-zero covariant curvature tensors fields of second order.

$$(1.34) \quad \mathcal{B}_m \mathcal{B}_l U_{jkh}^i = w_{lm} U_{jkh}^i + v_{lm} (\delta_k^i g_{jh} - \delta_h^i g_{jk}) - 2 \mu_l y^r \mathcal{B}_r (\delta_k^i C_{jhm} - \delta_h^i C_{jkm}) \quad , \quad \text{where}$$

$\mathcal{B}_m \mathcal{B}_l$ is Berwald's covariant differential operator of the second order with respect to x^l and x^m respectively, $a_{lm} = \mathcal{B}_m \lambda_l + \lambda_l \lambda_m$ and $b_{lm} = \mathcal{B}_m \mu_l + \lambda_l \mu_m$ are non-zero covariant tensors field and μ_l is non-zero covariant vector field of first order.

2. An Extended Framework for Generalized $\mathcal{B}_h$ U-Birecurrent Finsler Spaces

An advanced of generalized birecurrent structures in Finsler spaces related to the h(v)-curvature tensor is shown in this section. In the context of Berwald's connection, the goal of this section is to develop a generalized recurrence condition incorporating higher-order covariant derivatives.

Let F_n be an n-dimensional Finsler space with the Cartan connection. Let U_{jkh}^i denote be the space's h(v)-curvature tensor.

Assume that the generalized recurrent condition is satisfied by the h(v)-curvature tensor the following condition:

$$(2.1) \quad \mathcal{B}_l U_{jkh}^i = \lambda_l U_{jkh}^i + \mu_l (\delta_k^i g_{jh} - \delta_h^i g_{jk}) + \frac{1}{4} \eta_l (U_k^i g_{jh} - U_h^i g_{jk}) \quad , \quad U_{jkh}^i \neq 0 \quad ,$$

Where U_j^i is a deviation tensor connected to the curvature tensor U_{jkh}^i , and λ_l , μ_l and η_l are non-zero covariant vectors fields of first order.

The tensor U_{jkh}^i will be referred to as a generalized recurrent tensor, and a Finsler space that satisfies condition (2.1) will be called a generalized recurrent Finsler space. We will refer to the tensor as the generalization generalized $\mathcal{B}_h$ U-recurrent tensor and. We will abbreviate them as $G^{2nd} \mathcal{B}_h U-RF_n$ and $G^{2nd} \mathcal{B}_h U-R$, respectively.

Remark. 2.1. If we substitute any other deviation tensors, such as K_k^i and K_h^i or R_k^i and R_h^i or H_k^i and H_h^i , or N_k^i and N_h^i , for U_k^i and U_h^i , we obtain a new condition that characterizes a generalized generic $\mathcal{B}_h$ U-recurrent Finsler space in condition (2.1).

Taking $\mathcal{B}$ -covariant derivative for (2.1) with respect to x^m , in the since of Berwald and using (1.8) we get

$$(2.2) \quad \mathcal{B}_m \mathcal{B}_l U_{jkh}^i = (\mathcal{B}_m \lambda_l) U_{jkh}^i + \lambda_l (\mathcal{B}_m U_{jkh}^i) + (\mathcal{B}_m \mu_l) (\delta_j^i g_{kh} - \delta_k^i g_{jh})$$

$$-2 \mu_l y^r \mathcal{B}_r (\delta_k^i C_{jhm} - \delta_h^i C_{jkm}) + \frac{1}{4} (\mathcal{B}_m \eta_l) (U_k^i g_{jh} - U_h^i g_{jk}) \\ + \frac{1}{4} \eta_l [(\mathcal{B}_m U_k^i) g_{jh} - (\mathcal{B}_m U_h^i) g_{jk}] - \frac{1}{2} \eta_l y^r (U_k^i \mathcal{B}_r C_{jhm} - U_h^i \mathcal{B}_r C_{jkm}) .$$

Using (2.1) in (2.2), we get

$$(2.3) \quad \mathcal{B}_m \mathcal{B}_l U_{jkh}^i = w_{lm} U_{jkh}^i + v_{lm} (\delta_k^i g_{jh} - \delta_h^i g_{jk}) + \frac{1}{4} c_{lm} (U_k^i g_{jh} - U_h^i g_{jk}) \\ - 2 \mu_l y^r \mathcal{B}_r (\delta_k^i C_{jhm} - \delta_h^i C_{jkm}) - \frac{1}{2} \eta_l y^r (U_k^i \mathcal{B}_r C_{jhm} - U_h^i \mathcal{B}_r C_{jkm}) \\ + \frac{1}{4} \eta_l [(\mathcal{B}_m U_k^i) g_{jh} - (\mathcal{B}_m U_h^i) g_{jk}] .$$

Where $w_{lm} = \mathcal{B}_m \lambda_l + \lambda_l \lambda_m$, $v_{lm} = \lambda_l \mu_m + \mathcal{B}_m \mu_l$ and $c_{lm} = \lambda_l \eta_m + \mathcal{B}_m \eta_l$ are non-zero covariant tensors field of second order and η_l and μ_l are non-zero covariant vectors fields.

Definition 2.1. A Finsler space F_n for which $h(v)$ -curvature tensor U_{jkh}^i satisfies the condition (2.3), where w_{lm} , v_{lm} and c_{lm} are non-zero covariant tensors fields of second order, and η_l and μ_l are non-zero covariant vectors fields of first order, is called *generalization generalized $\mathcal{B}_h U$ -birecurrent Finsler Space* and the tensor will be called the *generalization generalized $\mathcal{B}U$ -recurrent tensor*, we shall denoted them briefly by $G^{2nd} \mathcal{B}_h U\text{-BRF}_n$ and $G^{2nd} \mathcal{B}U\text{-BRF}_n$ respectively.

Result 2.1. All $G^{2nd} \mathcal{B}_h U\text{-BRF}_n$ is $G^{2nd} \mathcal{B}U\text{-BRF}_n$.

Transvecting, (2.3) by y^h and using (1.15d), (1.8c), (1.2), (1.4b), (1.6a) and (1.6d), we get

$$(2.4) \quad \mathcal{B}_m \mathcal{B}_l U_{jk}^i = w_{lm} U_{jk}^i + v_{lm} (\delta_k^i y_j - y^i g_{jk}) + \frac{1}{4} c_{lm} (U_k^i y_j - U_h^i y^h g_{jk}) \\ + 2 \mu_l y^r \mathcal{B}_r (y^i C_{jkm}) + \frac{1}{2} \eta_l y^r (U_h^i y^h \mathcal{B}_r C_{jkm}) + \frac{1}{4} \eta_l [(\mathcal{B}_m U_k^i) y_j - (\mathcal{B}_m U_h^i y^h) g_{jk}] .$$

Suppose that $U_h^r y^h = U^r$ in (2.4) and using (1.8c), we get

$$(2.5) \quad \mathcal{B}_m \mathcal{B}_l U_{jk}^i = w_{lm} U_{jk}^i + v_{lm} (\delta_k^i y_j - y^i g_{jk}) + \frac{1}{4} c_{lm} (U_k^i y_j - U^i g_{jk}) \\ + 2 \mu_l y^r y^i \mathcal{B}_r C_{jkm} + \frac{1}{2} \eta_l y^r U^i \mathcal{B}_r C_{jkm} + \frac{1}{4} \eta_l [(\mathcal{B}_m U_k^i) y_j - (\mathcal{B}_m U^i) g_{jk}]$$

Contracting i and h in (2.3) and using (1.4g), (1.4c), (1.6a) and (1.17b), we get

$$(2.6) \quad \mathcal{B}_m \mathcal{B}_l U_{jk} = a_{lm} U_{jk} + (1-n) b_{lm} g_{jk} + \frac{1}{4} c_{lm} (U_k^s g_{js} - U_s^s g_{jk}) - 2 \mu_l y^r \mathcal{B}_r (\delta_k^s C_{jsm} - \delta_s^s C_{jkm}) \\ - \frac{1}{2} \eta_l y^r (U_k^s \mathcal{B}_r C_{jsm} - U_s^s \mathcal{B}_r C_{jkm}) + \frac{1}{4} \eta_l [(\mathcal{B}_m U_k^s) g_{js} - (\mathcal{B}_m U_s^s) g_{jk}] .$$

Suppose that $U_r^r = U$, $U_k^r g_{jr} = U_{jk}$ and $\delta_k^r C_{jrm} = C_{jkm}$ in the above equation, we get

$$(2.7) \quad \mathcal{B}_m \mathcal{B}_l U_{jk} = a_{lm} U_{jk} + (1-n) (b_{lm} g_{jk} - 2 \mu_l y^r \mathcal{B}_r C_{jkm}) + \frac{1}{4} c_{lm} (U_{jk} - U g_{jk}) \\ - \frac{1}{2} \eta_l y^r (U_k^s \mathcal{B}_r C_{jsm} - U \mathcal{B}_r C_{jkm}) + \frac{1}{4} \eta_l [(\mathcal{B}_m U_k^s) g_{js} - (\mathcal{B}_m U) g_{jk}] .$$

Thus, we may conclude

Theorem. 2.1. In $G^{2nd} \mathcal{B}_h U\text{-BRF}_n$ then, the $h(v)$ -torsion tensor U_{jk}^i and the Ricci tensor U_{jk} are given by (3.5) and (3.7), respectively.

Contracting i and k in (2.4), using (1.4c), (1.4a), (1.8c), (1.6a) and (1.17c), we get

$$(2.8) \quad \mathcal{B}_m \mathcal{B}_l C_{js}^s = w_{lm} C_{jk}^s + (n-1) y_j v_{lm} + \frac{1}{4} c_{lm} (U_s^s y_j - U^s g_{js}) + 2 \mu_l y^r \mathcal{B}_r (0) \\ + \frac{1}{2} \eta_l y^r (U^i \mathcal{B}_r C_{jkm}) + \frac{1}{4} \eta_l [(\mathcal{B}_m U_s^s) y_j - (\mathcal{B}_m U^s) g_{js}] .$$

Suppose that $U_s^s = U$ and $U^s g_{js} = U_j$, we get

$$(2.9) \quad \mathcal{B}_m \mathcal{B}_l C_{js}^s = w_{lm} C_{jk}^s + (n-1) y_j v_{lm} + \frac{1}{4} c_{lm} (U y_j - U_j) \\ + \frac{1}{2} \eta_l y^r (U^s \mathcal{B}_r C_{jsm}) + \frac{1}{4} \eta_l [(\mathcal{B}_m U) y_j - (\mathcal{B}_m U^s) g_{js}] .$$

Thus, we may conclude

Theorem. 2.2. In $G^{2nd} \mathcal{B}_h U\text{-BRF}_n$, the curvature tensor C_{js}^s satisfies (2.9).

3. Some Relationship among Curvatures Tensors In $G^{2nd} \mathcal{B}_h U\text{-BRF}_n$

Several relations between various curvature tensors in generalized birecurrent Finsler spaces are established in this section. Specifically, we examine how the Berwald curvature tensor, Cartan curvature tensors, and normal projective curvature tensor interacts.

By using Remark (2.1) in (2.1) and instead of U_k^i and U_h^i by the deviation tensor K_k^i and K_h^i , we get

$$(3.1) \quad \mathcal{B}_l U_{jkh}^i = \Lambda_l U_{jkh}^i + \mu_l (\delta_k^i g_{jh} - \delta_h^i g_{jk}) + \frac{1}{4} \eta_l (U_k^i g_{jh} - U_h^i g_{jk}), \quad U_{jkh}^i \neq 0.$$

Taking $\mathcal{B}$ -covariant derivative for (3.1) with respect to x^m , in the since of Berwald and using (1.9), we get

$$(3.2) \quad \begin{aligned} \mathcal{B}_m (\mathcal{B}_l U_{jkh}^i) &= (\mathcal{B}_m \lambda_l) U_{jkh}^i + \lambda_l (\mathcal{B}_m U_{jkh}^i) + (\mathcal{B}_m \mu_l) (\delta_k^i g_{jh} - \delta_h^i g_{jk}) \\ &- 2 \mu_l y^r \mathcal{B}_r (\delta_k^i C_{jhm} - \delta_h^i C_{jkm}) + \frac{1}{4} (\mathcal{B}_m \eta_l) (K_k^i g_{jh} - K_h^i g_{jk}) \\ &+ \frac{1}{4} \eta_l [(\mathcal{B}_m K_k^i) g_{jh} - (\mathcal{B}_m K_h^i) g_{jk}] - \frac{1}{2} \eta_l y^r \mathcal{B}_r [K_k^i (\mathcal{B}_r C_{jhm}) - K_h^i (\mathcal{B}_r C_{jkm})]. \end{aligned}$$

Using (3.1) in (3.2), we get

$$(3.3) \quad \begin{aligned} \mathcal{B}_m \mathcal{B}_l U_{jkh}^i &= w_{lm} U_{jkh}^i + v_{lm} (\delta_k^i g_{jh} - \delta_h^i g_{jk}) + \frac{1}{4} c_{lm} (K_k^i g_{jh} - K_h^i g_{jk}) \\ &- 2 \mu_l y^r \mathcal{B}_r (\delta_k^i C_{jhm} - \delta_h^i C_{jkm}) + \frac{1}{4} \eta_l [(\mathcal{B}_m K_k^i) g_{jh} - (\mathcal{B}_m K_h^i) g_{jk}] \\ &- \frac{1}{2} \eta_l y^r [K_k^i (\mathcal{B}_r C_{jhm}) - K_h^i (\mathcal{B}_r C_{jkm})]. \end{aligned}$$

Where $w_{lm} = \mathcal{B}_m \lambda_l + \lambda_l \lambda_m$, $v_{lm} = \lambda_l \mu_m + \mathcal{B}_m \mu_l$ and $c_{lm} = \lambda_l \eta_l + \mathcal{B}_m \eta_l$ are non-zero covariant tensors fields of second order, respectively.

Thus, we conclude

Theorem.3.3. A $G^{2nd} \mathcal{B}_h U\text{-}BRF_n$, may characterized by the condition (3.3).

By using (1.11) and (1.14d) in (3.3), we get

$$(3.5) \quad \begin{aligned} \mathcal{B}_m \mathcal{B}_l U_{jkh}^i &= w_{lm} U_{jkh}^i + \frac{1}{2} c_{lm} K_{jkh}^i + v_{lm} (\delta_k^i g_{jh} - \delta_h^i g_{jk}) + \frac{1}{4} \eta_l [(\mathcal{B}_m K_k^i) g_{jh} \\ &- (\mathcal{B}_m K_h^i) g_{jk}] - 2 \mu_l y^r \mathcal{B}_r (\delta_k^i C_{jhm} - \delta_h^i C_{jkm}) - \frac{1}{2} \eta_l y^r (K_k^i \mathcal{B}_r C_{jhm} - K_h^i \mathcal{B}_r C_{jkm}). \end{aligned}$$

Thus, we may conclude

Corollary 3.1. In $G^{2nd} \mathcal{B}_h U\text{-}BRF_n$, the relationship between the h(v)-curvature tensor U_{jkh}^i and Cartan's fourth curvature tensor K_{jkh}^i is given by (3.5).

By using (1.16) in (3.3), we get

$$(3.6) \quad \begin{aligned} \mathcal{B}_m \mathcal{B}_l U_{jkh}^i &= w_{lm} U_{jkh}^i + \frac{1}{4} c_{lm} H_{jkh}^i - \frac{1}{4} c_{lm} y^s (\partial_j K_{skh}^i) + v_{lm} (\delta_k^i g_{jh} - \delta_h^i g_{jk}) \\ &+ \frac{1}{4} \eta_l [(\mathcal{B}_m K_k^i) g_{jh} - (\mathcal{B}_m K_h^i) g_{jk}] - 2 \mu_l y^r \mathcal{B}_r (\delta_k^i C_{jhm} - \delta_h^i C_{jkm}) \\ &- \frac{1}{2} \eta_l y^r [K_k^i (\mathcal{B}_r C_{jhm}) - K_h^i (\mathcal{B}_r C_{jkm})]. \end{aligned}$$

Thus, we may conclude

Corollary 3.2. In $G^{2nd} \mathcal{B}_h U\text{-}BRF_n$, the relationship between n the h(v)-curvature tensor U_{jkh}^i and Berwald's curvature tensor H_{jkh}^i given by (3.6).

Let's look at a Finsler space F_n where the h(v) curvature tensor U_{jkh}^i meet the requirements listed below:

$$(3.7) \quad \mathcal{B}_l U_{jkh}^i = \Lambda_l U_{jkh}^i + \mu_l (\delta_k^i g_{jh} - \delta_h^i g_{jk}) + \frac{1}{4} \eta_l (H_k^i g_{jh} - H_h^i g_{jk}), \quad U_{jkh}^i \neq 0.$$

Taking $\mathcal{B}$ -covariant derivative for (3.7) with respect to x^m , in the since of Berwald and using (1.9) we get

$$(3.8) \quad \begin{aligned} \mathcal{B}_m (\mathcal{B}_l U_{jkh}^i) &= (\mathcal{B}_m \lambda_l) U_{jkh}^i + \lambda_l \mathcal{B}_m U_{jkh}^i + (\mathcal{B}_m \mu_l) (\delta_k^i g_{jh} - \delta_h^i g_{jk}) \\ &- 2 \mu_l y^r \mathcal{B}_r (\delta_k^i C_{jhm} - \delta_h^i C_{jkm}) + \frac{1}{4} (\mathcal{B}_m \eta_l) (H_k^i g_{jh} - H_h^i g_{jk}) \\ &+ \frac{1}{4} \eta_l [(\mathcal{B}_m H_k^i) g_{jh} - (\mathcal{B}_m H_h^i) g_{jk}] - \frac{1}{2} \eta_l y^r \mathcal{B}_r [H_k^i (\mathcal{B}_r C_{jhm}) - H_h^i (\mathcal{B}_r C_{jkm})]. \end{aligned}$$

Using (3.7) in (3.8), we get

$$(3.9) \quad \begin{aligned} \mathcal{B}_m \mathcal{B}_l U_{jkh}^i &= w_{lm} U_{jkh}^i + v_{lm} (\delta_k^i g_{jh} - \delta_h^i g_{jk}) + \frac{1}{4} c_{lm} (H_k^i g_{jh} - H_h^i g_{jk}) \\ &- 2 \mu_l y^r \mathcal{B}_r (\delta_k^i C_{jhm} - \delta_h^i C_{jkm}) + \frac{1}{4} \eta_l [(\mathcal{B}_m H_k^i) g_{jh} - (\mathcal{B}_m H_h^i) g_{jk}] \end{aligned}$$

$$-\frac{1}{2}\eta_l y^r (H_k^i \mathcal{B}_r C_{jhm} - H_h^i \mathcal{B}_r C_{jkm}) .$$

Where $w_{lm} = \mathcal{B}_m \lambda_l + \lambda_l \lambda_m$, $v_{lm} = \lambda_l \mu_m + \mathcal{B}_m \mu_l$ and $c_{lm} = \lambda_l \eta_l + \mathcal{B}_m \eta_l$ are non-zero covariant tensors fields of second order and μ_l and η_l are non-zero covariant vector fields of first order.

Thus, we conclude

Theorem 3.2. A $G^{2nd} \mathcal{B}_h U\text{-}BRF_n$, may characterized by the condition (3.9).

Suppose that $H_k^i g_{jh} = H_{jkh}^i$ in (3.9), we get

$$(3.10) \quad \mathcal{B}_m \mathcal{B}_l U_{jkh}^i = w_{lm} U_{jkh}^i + v_{lm} (\delta_k^i g_{jh} - \delta_h^i g_{jk}) + \frac{1}{4} c_{lm} (H_{jkh}^i - H_{jkh}^i) \\ - 2 \mu_l y^r \mathcal{B}_r (\delta_k^i C_{jhm} - \delta_h^i C_{jkm}) + \frac{1}{4} \eta_l [(\mathcal{B}_m H_k^i) g_{jh} - (\mathcal{B}_m H_h^i) g_{jk}] \\ - \frac{1}{2} \eta_l y^r (H_k^i \mathcal{B}_r C_{jhm} - H_h^i \mathcal{B}_r C_{jkm}) .$$

By using (1.21) in (3.9), we get

$$(3.11) \quad \mathcal{B}_m \mathcal{B}_l U_{jkh}^i = w_{lm} U_{jkh}^i + \frac{1}{2} c_{lm} H_{jkh}^i + v_{lm} (\delta_k^i g_{jh} - \delta_h^i g_{jk}) \\ - 2 \mu_l y^r \mathcal{B}_r (\delta_k^i C_{jhm} - \delta_h^i C_{jkm}) + \frac{1}{4} \eta_l [(\mathcal{B}_m H_k^i) g_{jh} - (\mathcal{B}_m H_h^i) g_{jk}] \\ - \frac{1}{2} \eta_l y^r (H_k^i \mathcal{B}_r C_{jhm} - H_h^i \mathcal{B}_r C_{jkm}) .$$

Thus, we conclude

Corollary 3.3. In $G^{2nd} \mathcal{B}_h U\text{-}BRF_n$, the relationship between the h(v)-curvature tensor U_{jkh}^i and Berwald's curvature tensor H_{jkh}^i may given by (3.11).

Let's look at a Finsler space F_n where the h(v) curvature tensor U_{jkh}^i meet the requirements listed below:

$$(3.12) \quad \mathcal{B}_l U_{jkh}^i = \lambda_l U_{jkh}^i + \mu_l (\delta_k^i g_{jh} - \delta_h^i g_{jk}) + \frac{1}{4} \eta_l (H_k^i g_{jh} - H_h^i g_{jk}) , \quad U_{jkh}^i \neq 0 .$$

Taking B-covariant derivative for (3.12) with respect to x^m , in the since of Berwald and using (1.9) we get

$$(3.13) \quad \mathcal{B}_m (\mathcal{B}_l U_{jkh}^i) = (\mathcal{B}_m \lambda_l) U_{jkh}^i + \lambda_l (\mathcal{B}_m U_{jkh}^i) + (\mathcal{B}_m \mu_l) (\delta_k^i g_{jh} - \delta_h^i g_{jk}) \\ - 2 \mu_l y^r \mathcal{B}_r (\delta_k^i C_{jhm} - \delta_h^i C_{jkm}) + \frac{1}{4} (\mathcal{B}_m \eta_l) (R_k^i g_{jh} - R_h^i g_{jk}) \\ + \frac{1}{4} \eta_l [(\mathcal{B}_m R_k^i) g_{jh} - (\mathcal{B}_m R_h^i) g_{jk}] - \frac{1}{2} \eta_l y^r \mathcal{B}_r (R_k^i \mathcal{B}_r C_{jhm} - R_h^i \mathcal{B}_r C_{jkm}) .$$

By using (3.12) in (3.13), we get

$$(3.14) \quad \mathcal{B}_m \mathcal{B}_l U_{jkh}^i = w_{lm} U_{jkh}^i + v_{lm} (\delta_k^i g_{jh} - \delta_h^i g_{jk}) + \frac{1}{4} c_{lm} (R_k^i g_{jh} - R_h^i g_{jk}) \\ - 2 \mu_l y^r \mathcal{B}_r (\delta_k^i C_{jhm} - \delta_h^i C_{jkm}) + \frac{1}{4} \eta_l [(\mathcal{B}_m R_k^i) g_{jh} - (\mathcal{B}_m R_h^i) g_{jk}] \\ - \frac{1}{2} \eta_l y^r (R_k^i \mathcal{B}_r C_{jhm} - R_h^i \mathcal{B}_r C_{jkm}) ,$$

Where $w_{lm} = \mathcal{B}_m \lambda_l + \lambda_l \lambda_m$, $v_{lm} = \lambda_l \mu_m + \mathcal{B}_m \mu_l$ and $c_{lm} = \lambda_l \eta_l + \mathcal{B}_m \eta_l$ are non-zero covariant tensors fields of second order.

Thus, we may conclude

Theorem 3.3. A $G^{2nd} \mathcal{B}_h U\text{-}BRF_n$, may characterized by the condition (3.14).

Let's look at a Finsler space F_n where h(v)-curvature tensor U_{jkh}^i meet the requirements listed below:

$$(3.15) \quad \mathcal{B}_l U_{jkh}^i = \lambda_l U_{jkh}^i + \mu_l (\delta_k^i g_{jh} - \delta_h^i g_{jk}) + \frac{1}{4} \eta_l (N_k^i g_{jh} - N_h^i g_{jk}) , \quad U_{jkh}^i \neq 0 .$$

Taking B-covariant derivative for (3.15) with respect to x^m , in the since of Berwald and using (1.9) we get

$$(4.16) \quad \mathcal{B}_m (\mathcal{B}_l U_{jkh}^i) = (\mathcal{B}_m \lambda_l) U_{jkh}^i + \lambda_l (\mathcal{B}_m U_{jkh}^i) + (\mathcal{B}_m \mu_l) (\delta_k^i g_{jh} - \delta_h^i g_{jk}) \\ - 2 \mu_l y^r \mathcal{B}_r (\delta_k^i C_{jhm} - \delta_h^i C_{jkm}) + \frac{1}{4} (\mathcal{B}_m \eta_l) (N_k^i g_{jh} - N_h^i g_{jk}) \\ + \frac{1}{4} \eta_l [(\mathcal{B}_m N_k^i) g_{jh} - (\mathcal{B}_m N_h^i) g_{jk}] - \frac{1}{2} \eta_l y^r \mathcal{B}_r (N_k^i \mathcal{B}_r C_{jhm} - N_h^i \mathcal{B}_r C_{jkm}) .$$

Using (3.15) in (3.16), we get

$$(3.17) \quad \mathcal{B}_m \mathcal{B}_l U_{jkh}^i = w_{lm} U_{jkh}^i + v_{lm} (\delta_k^i g_{jh} - \delta_h^i g_{jk}) + \frac{1}{4} c_{lm} (N_k^i g_{jh} - N_h^i g_{jk}) \\ - 2 \mu_l y^r \mathcal{B}_r (\delta_k^i C_{jhm} - \delta_h^i C_{jkm}) + \frac{1}{4} \eta_l [(\mathcal{B}_m N_k^i) g_{jh} - (\mathcal{B}_m N_h^i) g_{jk}]$$

$$-\frac{1}{2}\eta_l y^r (N_k^i \mathcal{B}_r C_{jhm} - N_h^i \mathcal{B}_r C_{jkm}) .$$

Where $w_{lm} = \mathcal{B}_m \lambda_l + \lambda_l \mu_m$, $v_{lm} = \lambda_l \mu_m + \mathcal{B}_m \mu_l$ and $c_{lm} = \lambda_l \eta_l + \mathcal{B}_m \eta_l$ are non-zero covariant tensors of second order, respectively.

Theorem 3.3. A $G^{2nd} \mathcal{B}_h U\text{-}BRF_n$, may characterized by the condition (3.17).

Suppose that, $N_k^i g_{jh} = N_{jkh}^i$ in (3.16), we get

$$(3.18) \quad \mathcal{B}_m \mathcal{B}_l U_{jkh}^i = w_{lm} U_{jkh}^i + v_{lm} (\delta_k^i g_{jh} - \delta_h^i g_{jk}) + \frac{1}{4} c_{lm} (N_{jkh}^i - N_{jkh}^i) \\ - 2 \mu_l y^r \mathcal{B}_r (\delta_k^i C_{jhm} - \delta_h^i C_{jkm}) + \frac{1}{4} \eta_l [(\mathcal{B}_m N_k^i) g_{jh} - (\mathcal{B}_m N_h^i) g_{jk}] \\ - \frac{1}{2} \eta_l y^r (N_k^i \mathcal{B}_r C_{jhm} - N_h^i \mathcal{B}_r C_{jkm}) .$$

By using (1.31) in (3.18), we get

$$(3.19) \quad \mathcal{B}_m \mathcal{B}_l U_{jkh}^i = w_{lm} U_{jkh}^i + \frac{1}{2} c_{lm} N_{jkh}^i + v_{lm} (\delta_k^i g_{jh} - \delta_h^i g_{jk}) \\ - 2 \mu_l y^r \mathcal{B}_r (\delta_k^i C_{jhm} - \delta_h^i C_{jkm}) + \frac{1}{4} \eta_l [(\mathcal{B}_m N_k^i) g_{jh} - (\mathcal{B}_m N_h^i) g_{jk}] \\ - \frac{1}{2} \eta_l y^r (N_k^i \mathcal{B}_r C_{jhm} - N_h^i \mathcal{B}_r C_{jkm}) .$$

Thus, we conclude

Theorem 3.5. A $G^{2nd} \mathcal{B}_h U\text{-}BRF_n$, the relationship between h(v)-curvature tensor U_{jkh}^i and the normal projective curvature tensor N_{jkh}^i is given by (3.19).

Contracting the indices i and h in (3.19), using (1.15b), (1.32), (1.4c) and (1.4g), we get

$$(3.20) \quad \mathcal{B}_m \mathcal{B}_l U_{jk} = w_{lm} U_{jk} + \frac{1}{2} c_{lm} N_{jk} + (1-n)(v_{lm} g_{jk} - 2 \mu_l y^r \mathcal{B}_r C_{jkm}) \\ + \frac{1}{4} \eta_l [(\mathcal{B}_m N_k^s) g_{js} - (\mathcal{B}_m N_s^s) g_{jk}] - \frac{1}{2} \eta_l y^r (N_k^s \mathcal{B}_r C_{jsm} - N_s^s \mathcal{B}_r C_{jkm}) .$$

Thus, we conclude

Corollary 3.4. The equation (3.20) holds in $G^{2nd} \mathcal{B}_h U\text{-}BRF_n$.

Conclusions

The introduction of generalized U-birecurrent Finsler spaces has significantly enriched the field of Finsler geometry. Our research has revealed several novel properties of these spaces and has expanded our understanding of their geometric structure. The results presented in this paper open up new avenues for further research.

Recommendations for Future Research

- Explore the physical applications of generalized U-birecurrent Finsler spaces.
- Investigate the relationship between generalized U-birecurrent Finsler spaces and other geometric structures, such as conformal Finsler geometry.
- Develop numerical methods for studying the properties of generalized U-birecurrent Finsler spaces.

References

1. Abu-Donia, H., Shenawy, S., & Abdehameed, A. (2020). The w^* -curvature tensor on relativistic space-times. *Kyungpook Mathematical Journal*, 60(1), 185–195. <https://doi.org/10.5666/KMJ.2020.60.1.185>
2. Ahsan, Z., & Ali, M. (2016). Curvature tensor for the spacetime of general relativity. *Palestine Journal of Mathematics*, 5(1), 1–15.
3. Ahsan, Z., & Ali, M. (2014). On some properties of the w-curvature tensor. *Palestine Journal of Mathematics*, 3(1), 61–69.
4. Al-Qashbari, A. M. A. (2020). On generalized curvature tensors P_{jkh}^i of second order in Finsler space. *University of Aden Journal of Natural and Applied Sciences*, 24(1), 171–176.
5. Al-Qashbari, A. M. A. (2020). Recurrence decompositions in Finsler space. *Journal of Mathematical Analysis and Modeling*, 1, 77–86.
6. Al-Qashbari, A. M. A. (2020). Some identities for generalized curvature tensors in B-recurrent Finsler space. *Journal of New Theory*, 32, 30–39. <https://doi.org/10.17350/HJNP.2020.178>

7. Al-Qashbari, A. M., Abdallah, A. A., & Ahmed, F. A. (2024). On generalized birecurrent Finsler space of mixed covariant derivatives in Cartan sense. *International Journal of Research Publication and Reviews*, 5(8), 2834–2840. <https://doi.org/10.55248/gengpi.5.0824.1843>
8. Al-Qashbari, A. M., Abdallah, A. A., & Al-ssallal, F. A. (2024). Recurrent Finsler structures with higher-order generalizations defined by special curvature tensors. *International Journal of Advanced Research in Science, Communication and Technology*, 4(1), 68–75. <https://doi.org/10.48175/IJARST-15185>
9. Al-Qashbari, A. M., Abdallah, A. A., & Baleedi, S. M. (2025). Berwald covariant derivative and Lie derivative of conharmonic curvature tensors in generalized fifth recurrent Finsler space. *GPH-International Journal of Mathematics*, 8(1), 24–32.
10. Al-Qashbari, A. M. A., Abdallah, A. A., & Baleedi, S. M. (2025). A study of M-projective curvature tensor in generalized recurrent Finsler space via Lie derivative. *International Journal of Applied Science and Mathematical Theory*, 11(1), 1–9.
11. Al-Qashbari, A. M., Haoues, M., & Al-ssallal, F. A. (2024). A decomposition analysis of Weyl's curvature tensor via Berwald's first and second-order derivatives in Finsler spaces. *Journal of Innovative Applied Mathematics and Computational Science*, 4(2), 201-213. <https://doi.org/10.46323/jiames.v4i2.215>
12. Al-Qashbari, A. M. A., & Hadi, W. H. A. (2025). Generalized trirecurrence of Cartan's second curvature tensor in P^h -recurrent spaces. *Journal of Science and Technology*, 30(10), 59–65.
13. Al-Qashbari, A. M. A., & Mubark, A. O. A. (2025). Decomposition of generalized recurrent tensor fields of RhR^h -nth order in Finsler manifolds. *Journal of Science and Technology*, 30(2), 99–105.
14. Al-Qashbari, A. M. A., & Qasem, M. M. (2026). Computational extensions of generalized R^h -recurrent Finsler spaces and applications to geometric machine learning. *Journal of Science and Technology*, 31(1), 50–57.
15. Al-Qashbari, A. M., Saleh, S., & Ibedou, I. (2024). On some relations of R-projective curvature tensor in recurrent Finsler space. *Journal of Non-Linear Modeling and Analysis*, 6(4), 1216–1227. <https://doi.org/10.12150/jnma.2024.1216>
16. [16] Al-Qashbari, A. M., et al. (2026). Investigating the relationships between Weyl's and Cartan's second curvature tensors in Finsler spaces. *International Journal of Analysis and Applications*, 24, 59. <https://doi.org/10.28924/2291-8639-24-59>
17. Al-Qashbari, A. M. A., & Husien, M. M. Q. (2025). Theoretical analysis of generalized W^h -birecurrent Finsler spaces with emphasis on Weyl's projective curvature. *Electronic Journal of University of Aden for Basic and Applied Sciences*, 6(4).
18. Hadi, W. H. A., Al-Qashbari, A. M. A., & Hanbala, A. M. A. (2025). Generalized H^h -recurrent Finsler spaces and their relations with Berwald and Cartan curvature tensors. *Journal of the Faculties of Education – University of Aden*, 19(2), 913–920.
19. Mandal, A., Yıldırım, M., Al-Qashbari, A. M. A., & Khan, M. N. I. (2025). Some solitons on anti-invariant submanifolds of para-Sasakian manifolds. *Journal of the Faculties of Education – University of Aden*, 19(2), 959–968.
20. Al-Qashbari, A. M. A., & Al-Arashi, W. H. (2025). Generalized-recurrent Finsler geometry with applications to anisotropic image processing. *Journal of Science & Technology*, 30(6).
21. Al-Qashbari, A. M. A., Abdallah, A. A., & Baleedi, S. M. (2025). Lie derivative approach to scalar quantities of tensors in generalized recurrent Finsler space. *Scholar Journal for Sciences & Technology*, 3(6).
22. Al-Qashbari, A. M. A., & Al-ssallal, F. A. M. (2025). Higher-order Cartan derivatives and curvature tensor decomposition in Finsler spaces. *Electronic Journal of University of Aden for Basic and Applied Sciences*, 6(2).
23. Al-Qashbari, A. M. A., Munassar, N., Abdullah, M. F., & Al-Hurdi, A. S. K. (2025). Enhancing performance through dynamic AES techniques. *Technological Applied and Humanitarian Academic Journal*, 2(1).
24. Al-Qashbari, A. M. A., Abdallah, A. A., & Baleedi, S. M. (2025). On concircular vector field and Lie derivative in generalized recurrent Finsler space. *Journal of Finsler Geometry and Its Applications*, 6(2), 82–92.

25. Al-Qashbari, A. M. A. (2025). Geometric properties and recurrence conditions in generalized H^v -recurrent Finsler spaces. *Journal of Education Faculties*, 19(1), 500–510.
26. Al-Qashbari, A. M. A., Abdallah, A. A., & Nasr, K. S. (2025). On generalized trirecurrent space using covariant derivatives in Finsler geometry. *Journal of Mathematical Problems, Equations and Statistics*, 6(1), 91–100.
27. Al-Qashbari, A. M. A., Abdallah, A. A., & Ahmed, F. A. (2025). Generalized tri-recurrent Finsler space under mixed covariant derivatives. *International Journal of Advanced Research in Science, Communication and Technology*, 2025.
28. Al-Qashbari, A. M. A., Abdallah, A. A., Baleedi, S. M., et al. (2025). The Kulkarni–Nomizu product in generalized recurrent Finsler spaces. *Journal of Applied and Pure Mathematics*, 7(3–4), 277–291.
29. Al-Qashbari, A. M. A., Abdallah, A. A., & Al-Ssallal, F. A. (2025). Curvature properties of generalized trirecurrent spaces in Finsler geometry. *International Journal of Research Publication and Reviews*, 6(6), 6341–6352.
30. Al-Qashbari, A. M. A., Abdallah, A. A., & Baleedi, S. M. (2025). Projective motions in generalized recurrent Finsler space. *International Journal of Research Publication and Reviews*, 6(4), 16101–16106.
31. Rund, H. (1959). *The Differential Geometry of Finsler Spaces*. Springer. <https://doi.org/10.1007/978-3-642-53099-9>
32. Bao, D., Robles, C., & Shen, Z. (2024). Advances in Finsler geometry and curvature structures. *Differential Geometry and Its Applications*, 92, 101846. <https://doi.org/10.1016/j.difgeo.2024.101846>
33. Cheng, X., & Shen, Z. (2024). Curvature properties of Finsler manifolds and geometric applications. *Journal of Geometry and Physics*, 198, 105102. <https://doi.org/10.1016/j.geomphys.2024.105102>
34. Deng, S., & Hou, Z. (2025). Curvature tensors and geometric invariants in modern Finsler geometry. *Results in Mathematics*, 80, 45. <https://doi.org/10.1007/s00025-025-01945-6>
35. Li, B., & Mo, X. (2024). Tensorial structures and curvature analysis in Finsler manifolds. *Mediterranean Journal of Mathematics*, 21, 112. <https://doi.org/10.1007/s00009-024-02412-7>
36. Shen, Z. (2024). Recent developments in Finsler geometry and curvature theory. *International Journal of Geometric Methods in Modern Physics*, 21(5), 2450072. <https://doi.org/10.1142/S0219887824500723>

فئة جديدة من فضاءات فنسلر المعممة ذات التكرار الثنائي-U وخصائص انحنائها

كمال سالم ناجي نصر¹ عادل محمد علي القشيري²

قسم الرياضيات، كلية التربية عدن - جامعة عدن - عدن، اليمن

² قسم الهندسة الطبية الحيوية، كلية الهندسة والحاسبات، جامعة العلوم والتكنولوجيا - عدن، اليمن

الملخص: في هذا البحث، نقدم فئة جديدة من فضاءات فنسلر المعممة ذات التكرار الثنائي-U، والمعرفة بدلالة مشتقات بيروالد التوافقية من الرتبة الثانية للتنبؤ الانحنائي $h(v)$. يوسع هذا التركيب الجديد المفاهيم القائمة لفضاءات فنسلر التكرارية وذات التكرار الثنائي، وذلك من خلال إدخال شروط تفاضلية من رتب أعلى. وقد تم إثبات عدة خصائص أساسية للفضاءات المقترحة، بما في ذلك العلاقات بين التنبؤ الانحنائي $h(v)$ وغيره من تنسورات الانحناء المهمة، مثل تنسورات انحناء كارتان، وتنسور انحناء بيروالد، وتنسور الانحناء الإسقاطي العادي. علاوة على ذلك، تم اشتقاق مجموعة من النظريات والنتائج التابعة لتوصيف السلوك الهندسي لهذه الفضاءات تحت شروط التكرار المعمم. وتوفر النتائج المتحصّل عليها إطارًا أوسع لدراسة البنى الانحنائية في هندسة فنسلر، كما تسهم في التطوير المستمر للتحليل الهندسي من الرتب العليا.

كلمات مفتاحية: هندسة فنسلر؛ الفضاء المعمم ذو التكرار الثنائي؛ التكرار الثنائي-U؛ مشتقة بيروالد التوافقية؛ التنبؤ الانحنائي؛ تنسور انحناء كارتان؛ تنسور الانحناء الإسقاطي.